## Summary of Clinical Preventive Services Guidelines for Adolescents up to Age 18 (CPSG-ADOL Summary) UCSF Division of Adolescent and Young Adult Medicine

NAHIC

Guidelines as of 10/2018, subject to change.

NAHIC			Guidelines as of 10/2018, subject to chang
<b>Preventive Services</b>	All $()$	At Risk (+)	Screening Test/ Procedure and Other Notes
Nutrition/exercise/obesity			
☐ Hypertension/Blood Pressure <sup>†</sup>	√		√ Bright Futures, USPSTF insufficient evidence
□ Obesity/BMI	V		Screen $\geq$ 6 years; offer/refer to appropriate intervention
☐ Cholesterol level		+	√ Bright Futures, USPSTF insufficient evidence
☐ Healthy diet <sup>†</sup>		+	√ Bright Futures, NHLBI
☐ Dyslipidemia	V		√ Bright Futures recommends one screening between ages 9-11 & 17-21,
			USPSTF insufficient evidence
Substance Use		T	
☐ Alcohol (screening and counseling) <sup>†</sup>	1		√ Bright Futures, USPSTF insufficient evidence
☐ Tobacco screening <sup>†</sup>	1		Provide interventions (education, brief counseling)
☐ Illicit Drugs (screening and counseling	(s) \ \ \		√ Bright Futures* and ACOG**, USPSTF insufficient evidence
Mental Health/Depression			
□ Depression (screening and treatmen	<b>t</b> ) √		Screen for MDD $\geq$ age 12, w/ adequate systems in place
□ Suicide Screening	V		√ Bright Futures and ACOG, USPSTF insufficient evidence
Safety/Violence			
□ Family/partner violence	1		Screen women of childbearing age
☐ Fighting	V		√ Bright Futures and ACOG
☐ Helmets	1		√ Bright Futures and ACOG
☐ Seat belts	1		√ Bright Futures and ACOG
□ Guns	1		√ Bright Futures and ACOG
□ Bullying	V		√ Bright Futures only
Reproductive Health			
□ HIV <sup>†</sup>	1	+	Bright Futures and USPSTF recommend one screening between ages 15-18, and annually for those at increased risk
☐ STI (screening and counseling) <sup>†</sup>		+	High-Intensity Counseling Interventions
□ Syphilis		+	VDRL
☐ Gonorrhea (females)		+	NAATs; test if ≤ 24 and sexually active
☐ Chlamydia (females)		+	NAATs; test if ≤ 24 and sexually active
☐ Chlamydia & Gonorrhea (male)		+	√ Bright Futures only
☐ Birth Control Methods	V	+	√ ACOG, + Bright Futures
□ Pregnancy		+	+ Bright Futures
Cancer Screening	l I	·	
□ Skin Cancer (counseling) <sup>†</sup>		+	Counsel those with fair skin ages 10-24 about reducing UV exposu
□ BRCA-Related Cancer <sup>†</sup>		+	Family Hx of breast, ovarian, tubal, or peritoneal cancer
Infectious Diseases including CDC Im	munization		
□ Td/Tdap	1		Td booster every 10 years
☐ Human papillomavirus	1		9vHPV vaccine for males and females up to age 26; 3 lifetime dose
□ Varicella (LIVE VACCINE)	√***		2 lifetime doses at least 4 weeks apart ***See below
☐ Measles, mumps, rubella	1		1 or 2 lifetime doses at least 4 weeks apart
☐ Influenza	N 1		1 dose annually
□ Pneumococcal	V	+	PCV13: 1 lifetime dose   PPSV23: 1-2 lifetime doses
☐ Hepatitis A	V		2 or 3 lifetime doses
☐ Hepatitis B	V		3 lifetime doses
☐ Meningococcal Quadrivalent	V		2 lifetime doses
□ Serogroup B Meningococcal	,	+	Men B vaccine (2 or 3-dose series) to those 16-23 years old
÷		+	Anti–HCV antibody testing, polymerase chain reaction testing
110punius o gerteming		mmandations are f	220 , unitarity totaling, polymeruse chain reaction testing

**Bold** = US Preventive Services Task Force (USPSTF) A or B Recommendation or CDC recommendations for immunizations.

Current evidence is insufficient to assess the balance of benefits and harms of service.

 $\sqrt{\ }$  = All adolescents + = Adolescents at risk For more information, please view the appendix, and

visit the official website.

Cite as: National Adolescent and Young Adult Health Information Center (2018). Summary of Clinical Preventive Services Guidelines for Adolescents up to Age 18. San Francisco, CA: National Adoelscent and Young Adult Health Information Center, University of California, San Francisco. Retrieved from: <a href="http://nahic.ucsf.edu/resource\_center/adolescent-guidelines/">http://nahic.ucsf.edu/resource\_center/adolescent-guidelines/</a>.

<sup>\*</sup> Bright Futures: recommendations are for annual visits, up to age 21.

<sup>\*\*</sup> American Congress of Obstetricians and Gynecologists (ACOG) recommendations, up to age 26.

<sup>\*\*\*</sup> The varicella vaccine should NOT be given to patients with these contraindications.

<sup>†</sup> USPSTF update in progress.